

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
25 August 2005 (25.08.2005)

PCT

(10) International Publication Number  
**WO 2005/077881 A1**

(51) International Patent Classification<sup>7</sup>: **C07C 69/618**,  
C11D 3/50, A61K 7/46

(21) International Application Number:  
PCT/CH2005/000074

(22) International Filing Date: 10 February 2005 (10.02.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0403115.9 12 February 2004 (12.02.2004) GB

(71) Applicant (for all designated States except US): GIVAUDAN SA [CH/CH]; Chemin de la Parfumerie 5, CH-1214 Vernier (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): FLACHSMANN, Felix [CH/CH]; Langhagweg 2, CH-8600 Duebendorf (CH). BACHMANN, Jean-Pierre [CH/CH]; Holzmoosrütisteig 4, CH-8820 Wädenswil (CH).

(74) Agent: MCSTEAL, John, Anthony; Ueberlandstrasse 138, CH-8600 Duebendorf (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:


— as to the identity of the inventor (Rule 4.17(i)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ.

[Continued on next page]

(54) Title: 3-(2-ALKOXYCARBONYLOXY-PHENYL) ACRYLIC ACID ESTERS AND THEIR USE AS PRECURSORS FOR THE DELIVERY OF OLFACTORY COMPOUNDS



Solution D



(57) Abstract: A compound of formula (I), their use as precursors and a method of their production wherein n, Y, R, R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup> has the same meaning as given in the specification.

WO 2005/077881 A1



NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA,

ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— of inventorship (Rule 4.17(iv)) for US only

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

3-(2-ALKOXYCARBONYLOXY-PHENYL) ACRYLIC ACID ESTERS AND THEIR USE AS PRECURSORS  
FOR THE DELIVERY OF OLFACTORY COMPOUNDS


The present invention refers to 3-(2-alkoxycarbonyloxy-phenyl)acrylic acid esters and their use as precursors for the delivery of olfactory compounds. This invention relates  
5 furthermore to a method of their production and to consumer products comprising them.

A principal strategy currently employed in imparting odors to consumer products is the admixing of the fragrance directly into the product. There are, however, several drawbacks to this strategy. The fragrance material can be too volatile and/or too soluble  
10 in water, resulting in fragrance loss during manufacturing, storage, and use. Many fragrance materials are also unstable over time. This again results in loss during storage. In many consumer products it is desirable for the fragrance to be released slowly over time. Microencapsulation and inclusion complexes with cyclodextrins have been used to help decrease volatility, improve stability and provide slow-release  
15 properties. However, these methods are for a number of reasons often not successful. In addition, cyclodextrins can be too expensive for use in many applications. It is therefore desirable to have a fragrance delivery system which is capable of releasing the fragrant compound or compounds in a controlled manner, maintaining a desired odour or fragrance over a prolonged period of time.

20

The principle of using precursors for the delivery of fragrance compounds appeared for the first time some years ago in the literature. The use of 3-(2-hydroxyaryl)acrylic acid esters (compound A below) is described in EP 0 936 211. This delivery system releases one or more olfactory compounds upon exposure to light. Using this system in various  
25 consumer products leads to a prolonged release of the fragrant compound(s). Unfortunately, the use of 3-(2-hydroxyaryl)acrylic acid esters may lead to discoloration, such as yellow discoloration, not only of consumer products, such as laundry care products, e.g. fabric softeners and detergents, comprising it, but it also may lead to discoloration of the substrate, for example, the fabric to which the product is applied  
30 during the washing cycle or rinse cycle. Discoloration of a product such as a fabric in general is not desired, thus there still remains a need for precursors having the advantageous ability to release one or more olfactory compounds, but without causing discoloration.

It has now been found that certain 3-(2-alkoxycarbonyloxy-phenyl)acrylic acid esters have the ability to release one or more olfactory compounds without showing discoloration to be visible to the naked eye. Surprisingly, it has been found that certain phenol protecting groups, in particular esters and carbonates, have the ability of rendering the prior art compounds (A)




A

color-stable in consumer products comprising them.

Thus, it is believed, without restricting the invention in any way, that the free phenolic hydroxyl group of the prior art compounds, as described for example in EP 0 936 211 and WO 03/022978, is responsible for the discoloration.

Accordingly, a first aspect of the present invention refers to the use of a compound of formula (I) as precursor for olfactory compounds



(I)

wherein the acrylic acid ester double bond is of the E configuration;

n is zero or 1;

Y is  $-CR^5R^6R^7$ , wherein  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen or a  $C_1$ -  $C_{18}$ , preferably  $C_1$ -  $C_{10}$ , hydrocarbon residue of which preferably at least one residue  $R^5$ ,  $R^6$

and  $R^7$  is not hydrogen, and the sum of all carbon atoms ( $R^5 + R^6 + R^7$ ) is not greater than 18, preferably the sum of all carbon atoms is between 6 and 15; or

Y is  $-CR^5R^6R^7$ , wherein  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen or a  $C_1$ - $C_{18}$ , preferably  $C_1$ - $C_{10}$ , hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), of which preferably at least one residue  $R^5$ ,  $R^6$  and  $R^7$  is not hydrogen, and the sum of all carbon atoms ( $R^5 + R^6 + R^7$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15, for example Y is 2-(2-butoxy-ethoxy)-ethyl; or

Y is  $-CR^8=CR^9R^{10}$ , wherein  $R^8$ ,  $R^9$  and  $R^{10}$  are independently hydrogen or a  $C_1$ - $C_{18}$ , preferably  $C_1$ - $C_{10}$ , hydrocarbon residue, of which preferably at least one of the residues  $R^8$ ,  $R^9$  and  $R^{10}$  is not hydrogen, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^8 + R^9 + R^{10}$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15; or

Y is  $-CR^8=CR^9R^{10}$ , wherein  $R^8$ ,  $R^9$  and  $R^{10}$  are independently hydrogen or a  $C_1$ - $C_{18}$ , preferably  $C_1$ - $C_{10}$ , hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), of which preferably at least one of the residues  $R^8$ ,  $R^9$  and  $R^{10}$  is not hydrogen, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^8 + R^9 + R^{10}$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15;


$R^2$  and  $R^3$  are independently hydrogen;  $C_1$ - $C_6$  alkyl, e.g. methyl, ethyl, iso-propyl, n-butyl, tert-butyl;  $C_1$ - $C_6$  alkoxy residue, e.g. methoxy, ethoxy;  $-NO_2$ ;  $-NH_2$ ;  $-NHCO_2CH_3$ ;  $-N(C_1-C_6 \text{ alkyl})_2$ , e.g. dimethylamino, diethylamino;  $-N(\text{hydroxyalkyl})_2$ , e.g. di(hydroxyethyl)amino, di(hydroxypropyl)amino;  $-NHC(O)-(C_1-C_8 \text{ alkyl})$  or  $-NHC(O)-(C_3-C_8 \text{ aryl})$ , e.g.  $-NHC(O)$ -methyl or  $-NHC(O)$ -phenyl; or

$R^2$  and  $R^3$  are attached at the positions C(6,7), C(7,8), or C(8,9), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

$R^4$  in 2- or 3-position is hydrogen;  $C_1$ - $C_4$  alkyl, e.g. methyl, ethyl, tert-butyl;  $C_2$ - $C_4$  alkenyl, e.g. vinyl, propenyl;  $C_3$ - $C_6$  cycloalkyl, e.g. cyclopropyl, cyclopentyl, cyclohexyl; or  $-CN$ ; and

if n is zero, R is a C<sub>1</sub>-C<sub>24</sub>, preferably C<sub>1</sub>-C<sub>18</sub>, hydrocarbon residue, e.g. methyl, ethyl or phenyl; or C<sub>1</sub>-C<sub>24</sub>, preferably C<sub>1</sub>-C<sub>18</sub>, hydrocarbon residue containing one or more heteroatoms selected from N, O and Si; or

if n is 1, R is a C<sub>1</sub>-C<sub>25</sub>, preferably C<sub>1</sub>-C<sub>18</sub>, hydrocarbon residue; a C<sub>1</sub>-C<sub>25</sub> hydrocarbon residue containing one or more atoms/groups selected from N, O, Si, and C(O); or C<sub>1</sub>-C<sub>25</sub>, preferably C<sub>1</sub>-C<sub>18</sub>, hydrocarbon residue substituted by an ionic substituent of the formula N(R<sup>20</sup>)<sub>3</sub><sup>+</sup>, in which R<sup>20</sup> is the residue of an alkyl group with 1 to 18 carbon atoms, preferably 1 to 8 carbon atoms, such as trimethylammonium, or tributylammonium; or R is a monovalent residue of the formula (i)



wherein

X is -CR<sup>14</sup>R<sup>15</sup>R<sup>16</sup>, wherein R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> are independently hydrogen or a C<sub>1</sub>-C<sub>18</sub>, preferably C<sub>1</sub>-C<sub>10</sub>, hydrocarbon residue, of which preferably at least one residue R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> is not hydrogen, and the sum of all carbon atoms (R<sup>14</sup>+R<sup>15</sup>+R<sup>16</sup>) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15; or

X is -CR<sup>14</sup>R<sup>15</sup>R<sup>16</sup>, wherein R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> are independently hydrogen or a C<sub>1</sub>-C<sub>18</sub>, preferably C<sub>1</sub>-C<sub>10</sub>, hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), of which preferably at least one residue R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> is not hydrogen, and the sum of all carbon atoms (R<sup>14</sup>+R<sup>15</sup>+R<sup>16</sup>) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15, for example X is 2-(2-butoxy-ethoxy)-ethyl; or

X is -CR<sup>17</sup>=CR<sup>18</sup>R<sup>19</sup>, wherein R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> are independently hydrogen or a C<sub>1</sub>-C<sub>18</sub>, preferably C<sub>1</sub>-C<sub>10</sub>, hydrocarbon residue, of which preferably at least one of the

residues  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  is not hydrogen, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^{17} + R^{18} + R^{19}$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15; or

5 X is  $-CR^{17}=CR^{18}R^{19}$ , wherein  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  are independently hydrogen or a  $C_1$ - $C_{18}$ , preferably  $C_1$ - $C_{10}$ , hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), of which preferably at least one of the residues  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  is not hydrogen, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^{17} + R^{18} + R^{19}$ ) is not greater than 18, preferable the  
10 sum of all carbon atoms is between 6 and 15;

$R^{12}$  and  $R^{13}$  are independently hydrogen;  $C_1$ - $C_6$  alkyl, e.g. methyl, ethyl, iso-propyl, n-butyl, tert-butyl;  $C_1$ - $C_6$  alkoxy residue, e.g. methoxy, ethoxy;  $-NO_2$ ;  $-NH_2$ ;  $-NHCO_2CH_3$ ;  $-N(C_1-C_6 \text{ alkyl})_2$ , e.g. dimethylamino, diethylamino;  $N(\text{hydroxyalkyl})_2$ , e.g. di(hydroxyethyl)amino, di(hydroxypropyl)amino;  $-NHC(O)-(C_1-C_8 \text{ alkyl})$ ; or  
15  $-NHC(O)-(C_3-C_8 \text{ aryl})$ , e.g.  $-NHC(O)$ -methyl or  $-NHC(O)$ -phenyl; or

$R^{12}$  and  $R^{13}$  are attached at the positions C(vi,vii), C(vii,viii), or C(viii,ix), and form together with the carbon atoms to which they are attached a dioxolane ring or a  
20 dioxane ring;

$R^{11}$  in ii- or iii-position is hydrogen;  $C_1$ - $C_4$  alkyl, e.g. methyl, ethyl, tert-butyl;  $C_2$ - $C_4$  alkenyl, e.g. vinyl, propenyl;  $C_3$ - $C_6$  cycloalkyl, e.g. cyclopropyl, cyclopentyl, cyclohexyl; or  $-CN$ .

25 As used in relation to the compounds of formula (I) "hydrocarbon residue" unless otherwise indicated refers to alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl alkylcycloalkyl, alkenylcycloalkyl, alkenylcycloalkenyl, aryl, alkylaryl or arylalkyl, and "hydrocarbon residues containing one or more atoms/groups selected from O, N and  
30 C(O)," refers to alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkenylcycloalkyl, alkenylcycloalkenyl, aryl, alkylaryl or arylalkyl wherein one or more carbon atoms are replaced by O, N and/or C(O).



As used herein by "olfactory compound" is meant a molecule having an odour, preferably a pleasant odour, detectable by a human. As used herein, the terms "olfactory" and "fragrant" are used interchangeably, and refer to the same compounds.

- 5 Compounds of formula (I) are preferred wherein n is 1, and Y is the residue of a fragrant alcohol  $\text{HO}-\text{CR}^5\text{R}^6\text{R}^7$  or the residue of the enol form of a fragrant aldehyde of the formula  $\text{O}=(\text{CH})-\text{CHR}^9\text{R}^{10}$ , or the residue of the enol form of a fragrant ketone of the formula  $\text{O}=(\text{CR}^8)-\text{CHR}^9\text{R}^{10}$  and if R is the monovalent residue of formula (i), X is the residue of a fragrant alcohol  $\text{HO}-\text{CR}^{14}\text{R}^{15}\text{R}^{16}$  or the residue of the enol form of a
- 10 fragrant aldehyde of the formula  $\text{O}=(\text{CH})-\text{CHR}^{18}\text{R}^{19}$  or the residue of the enol form of a fragrant ketone of the formula  $\text{O}=(\text{CR}^{17})-\text{CHR}^{18}\text{R}^{19}$ .

- Even more preferred are compounds of the present invention wherein n is 1, Y is the residue of a fragrant alcohol  $\text{HO}-\text{CR}^5\text{R}^6\text{R}^7$  or the residue of the enol form of a fragrant
- 15 aldehyde of the formula  $\text{O}=(\text{CH})-\text{CHR}^9\text{R}^{10}$  or the residue of the enol form of a fragrant ketone of the formula  $\text{O}=(\text{CR}^8)-\text{CHR}^9\text{R}^{10}$ , and R is selected from methyl, ethyl, propyl, butyl, pentyl, 2-ethylhexyl, cyclopentyl, cyclohexyl or the residue of a fragrant alcohol.

- For the purpose of the present invention the term "fragrant alcohol" is defined herein as
- 20 any alcohol having a molecular weight between 46 and 400, preferably between 100 and 300.

- Examples of fragrant alcohols of the formula  $\text{HO}-\text{CR}^5\text{R}^6\text{R}^7$  and  $\text{HO}-\text{CR}^{14}\text{R}^{15}\text{R}^{16}$  include: amyl alcohol; hexyl alcohol\*; 2-hexyl alcohol\*; heptyl alcohol\*; octyl alcohol\*; nonyl alcohol\*; decyl alcohol\*; undecyl alcohol\*; lauryl alcohol\*; myristic alcohol; 3-methyl-but-
- 25 2-en-1-ol\*; 3-methyl-1-pentanol; cis-3-hexenol\*; cis-4-hexenol\*; 3,5,5-trimethyl-hexanol; 3,4,5,6,6-pentamethylheptan-2-ol\*; citronellol\*; geraniol\*; oct-1-en-3-ol; 2,5,7-trimethyl-octan-3-ol; 2-cis-3,7-dimethyl-2,6-octadien-1-ol; 6-ethyl-3-methyl-5-octen-1-ol\*; 3,7-dimethyl-oct-3,6-dienol\*; 3,7-dimethyloctanol\*; 7-methoxy-3,7-dimethyl-octan-2-ol\*; cis-6-nonenol\*; 5-ethyl-2-nonanol; 6,8-dimethyl-2-nonanol\*; 2,2,8-trimethyl-7(8)-nonene-3-
- 30 ol; nona-2,6-dien-1-ol; 4-methyl-3-decen-5-ol\*; dec-9-en-1-ol \*\*; benzylalcohol; 2-methyl-undecanol; 10-undecen-1-ol; 1-phenyl-ethanol\*; 2-phenyl-ethanol\*; 2-methyl-3-phenyl-3-propenol; 2-phenyl-propanol\*; 3-phenyl-propanol\*; 4-phenyl-2-butanol; 2-methyl-5-phenyl-pentanol\*; 2-methyl-4-phenyl-pentanol\*; 3-methyl-5-phenyl-pentanol\*;
- 35 2-(2-methylphenyl)ethanol\*; 4-(1-methylethyl)benzene-methanol; 4-(4-hydroxyphenyl)-butan-2-one\*; 2-phenoxy-ethanol\*; 4-(1-methylethyl)-2-hydroxy-1-methyl benzene; 2-



methoxy-4-methyl-phenol; 4-methyl-phenol; anisic alcohol\*; p-tolyl alcohol\*; cinnamic alcohol\*; vanillin\*; ethyl vanillin\*; eugenol\*; isoeugenol\*; thymol; anethol\*; decahydro-2-naphthalenol; borneol\*; cedrenol\*; farnesol\*; fenchyl alcohol\*; menthol\*; 3,7,11-trimethyl-2,6,10-dodecatrien-1-ol; alpha ionol\*; tetrahydro ionol\*; 2-(1,1-dimethylethyl)cyclohexanol\*; 3-(1,1-dimethylethyl)cyclohexanol\*; 4-(1,1-dimethylethyl)cyclohexanol\*; 4-isopropyl-cyclohexanol; 6,6-dimethyl-bicyclo[3.3.1]hept-2-ene-2-ethanol; 6,6-dimethyl-bicyclo[3.1.1]hept-2-ene-methanol\*; p-menth-8-en-3-ol\*; 3,3,5-trimethyl-cyclohexanol; 2,4,6-trimethyl-3-cyclohexenyl-methanol\*; 4-(1-methylethyl)cyclohexyl-methanol\*; 4-(1,1-dimethylethyl)cyclohexanol; 2-(1,1-dimethylethyl)cyclohexanol; 2,2,6-trimethyl-alpha-propyl-cyclohexane propanol\*; 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol\*; 3-methyl-5-(2,2,3-trimethylcyclopentyl-3-enyl)pent-4-en-2-ol\*; 2-ethyl-4-(2,2,3-trimethylcyclopentyl-3-enyl)but-2-en-1-ol\*; 4-(5,5,6-trimethylbicyclo[2.2.1]hept-2-yl)cyclohexanol\*; 2-(2-methylpropyl)-4-hydroxy-4-methyl-tetrahydropyran\*; 2-cyclohexyl-propanol\*; 2-(1,1-dimethylethyl)-4-methyl-cyclohexanol\*; 1-(2-tert-butyl-cyclohexyloxy)-2-butanol\*; 1-(4-isopropyl-cyclohexyl)ethanol\*; 2,6-dimethyl-oct-7-en-2-ol\*\*; 2,6-dimethyl-heptan-2-ol\*\*; and 3,7-dimethyl-octa-1,6-dien-3-ol\*\*;

whereby \* indicates the preferred alcohols and \*\* indicate the more preferred alcohols.

For the purpose of the present invention the term "fragrant aldehyde" is defined herein as any aldehyde having a molecular weight between 100 and 450, preferably between 120 and 300.

Examples of fragrant aldehydes of the formula  $O=(CH)-CHR^9R^{10}$  and  $O=(CH)-CHR^{18}R^{19}$  include:

2,6,10-trimethylundec-9-enal\*; 1,2,3,4,5,6,7,8,-octahydro-8,8-dimethyl-2-naphthalenecarboxaldehyde; tridecanal; 2-[4-(1-methylethyl)phenyl]ethanal; 2,4-dimethyl-cyclohex-3-ene-1-carboxaldehyde\*; 4-carboxaldehyde-1,3,5-trimethyl-cyclohex-1-ene\*; 1-carboxaldehyde-2,4-dimethyl-cyclohex-3-ene\*; 1-carboxaldehyde-4-(4-hydroxy-4-methylpentyl)cyclohex-3-ene\*; 3,5,5-trimethyl-hexanal; heptanal\*; 2,6-dimethyl-hept-5-enal\*; decanal\*\*; dec-9-enal; dec-4-enal; 2-methyldecanal\*; undec-10-enal\*\*;

undecanal\*; dodecanal\*\*;

2-methyl-undecanal\*\*;

tridecanal; octanal\*\*;

nonanal\*; 3,5,5-trimethylhexanal; undec-9-enal\*\*;

2-phenyl-propanal\*; 4-methyl-phenyl-acetaldehyde\*;

3,7-dimethyl-octanal\*; dihydrofarnesal\*\*;

7-hydroxy-3,7-dimethyl-octanal\*; 2,6-dimethyl-oct-5-enal; 2-[4-(1-methylethyl)phenyl]ethanal\*;

3-(3-isopropyl-phenyl)butanal\*\*;

2-(3,7-dimethyloct-6-enoxy)ethanal; 1-carboxaldehyde-4-(4-methyl-3-pentenyl)cyclohex-3-

ene\* ; 2,3,5,5,-tetramethyl-hexanal; longifolic aldehyde; 2-methyl-4-(2,6,6-trimethylcyclohex-2-en-1-yl)butanal\*; 2-methyl-3-(4-tert-butylphenyl)propanal\*\*; 4-(1,1-dimethyl-ethyl)benzene-propanal\*; 2-[4-(1-methyl-ethyl)-phenyl]propanal; alpha-methyl-1,3-benzodioxole-5-propanal\*; 3,7-dimethyl-oct-6-enal\*; 2-methyl-3-(4-isopropylphenyl)-propionaldehyde\*; 4-(4-hydroxy-4-methyl-pentyl)cyclohex-3-en-1-carboxaldehyde\*\*;  
 5 alpha-methyl-1,3-benzodioxole-5-propanal\*; 1-carboxaldehyde-4-(1,1-dimethylethyl)-cyclohexane; 4-(octahydro-4,7-methano-5H-inden-5-ylidene)butanal; and [(3,7-dimethyl-6-octenyl)-oxy]acetaldehyde\*\*;  
 whereby \* indicates the preferred aldehydes and \*\* indicate the more preferred  
 10 aldehydes.

For the purpose of the present invention the term "fragrant ketone" is defined herein as any ketone having a molecular weight between 100 and 450, preferably between 120 and 350.

15 Examples of fragrant ketones of the formula  $O=(CR^8)-CHR^9R^{10}$  and  $O=(CR^{17})-CHR^{18}R^{19}$  include:

2-heptyl-cyclopentanone; 2,2,6,10-tetramethyltricyclo-[5.4.0.0(6,10)]undecan-4-one benzylacetone\*; carvone\*; 1,2,3,5,6,7-hexahydro-1,1,2,3,3,-pentamethyl-4H-inden-4-one\*; methyl heptenone\*; dimethyl octenone\*; 2-(butan-2-yl)cyclohexanone\*; 2-hexyl-  
 20 cyclopent-2-en-1-one\*; 2-(1-methylethyl)-5-methyl-cyclohexanone\*; 2-(2-methylethyl)-5-methyl-cyclohexanone\*; 3-methyl-cyclopentadecanone; 4-tert-pentyl-cyclohexanone\*; 3-oxo-2-pentyl-cyclopentane-acetic acid methyl ester\*\*; 1-(1,2,3,4,5,6,7,8-octahydro-2,3,8,8-tetramethyl-2-naphthalenyl)ethanone\*; and 3-methyl-5-propyl-cyclohex-2-en-1-one\*;

25 whereby \* indicates the preferred ketones and \*\* indicate the more preferred ketone.

Another embodiment relates to compounds of formula (I) wherein n is 1 and R and Y have the same meaning as given above and

- I)  $R^2$ ,  $R^3$  and  $R^4$  are H;
- 30 II)  $R^2$  and  $R^3$  is H, and  $R^4$  is methyl or -CN at C(2) or C(3), or phenyl at C(3);
- III)  $R^2$  is H,  $R^3$  is methyl, ethyl, propyl, or isopropyl at either C(6) to C(8) or methoxy, ethoxy, propyloxy at either C(6) to C(8), and  $R^4$  is H, methyl or -CN at C(2) or C(3), or phenyl at C(3);


IV)  $R^2$  and  $R^3$  is methyl at positions C(6,7), C(6,8), C(6,9), C(7,8), or C(8,9); or  $R^2$  and  $R^3$  is methoxy at C(7,9), and  $R^4$  is H, methyl or -CN at C(2) or C(3), or phenyl at C(3);

V)  $R^4$  is H, methyl or -CN at C(2) or C(3), or phenyl at C(3), and  $R^2$  is methyl at C(6) and  $R^3$  is isopropyl at C(9), or  $R^2$  is isopropyl at C(6) and  $R^3$  is methyl at C(9).

Compounds of formula (I) are preferred wherein n is 1, R and Y have the same meaning as given above,  $R^4$  is hydrogen or methyl at position C(2) or C(3), and  $R^2$  and  $R^3$  is hydrogen, or  $R^2$  is hydrogen and  $R^3$  is 7-methoxy, or  $R^2$  is hydrogen and  $R^3$  is 6-methyl, or  $R^2$  is hydrogen and  $R^3$  is 7-methyl, or  $R^2$  is hydrogen and  $R^3$  is 8-methyl, or  $R^2$  is hydrogen and  $R^3$  is 6-tert-butyl, or  $R^2$  is 6-tert-butyl and  $R^3$  is 8-tert-butyl.

The release of the active substances occurs in two successive steps via hydrolysis, preferably in the presence of enzymes, followed by photoisomerisation / lactonization, as shown in Scheme 1. Due to the two different consecutive cleavage mechanisms, it is possible to control the release of an olfactory compound at two different stages of the drying process of a substrate to which compounds of the present invention are applied. That is, for example, the release of an alcohol of the formula ROH (IV) after spin-drying of a fabric in a first step and the release of an alcohol (III), or for  $Y = -CR^8=CR^9R^{10}$  a ketone or aldehyde by tautomerisation, and a coumarin of formula (IIa), when exposed to UV light, e.g. sunlight during line-drying. In other words, a first boost of fragrance is perceivable when the washing machine is opened and a second boost of fragrance is perceivable during the line-drying process, if exposed to UV-light.

Scheme 1:




Accordingly, another aspect of the present invention is a process of providing an olfactory compound to a substrate comprising the steps:

- a) cleaving a compound of formula (I) by hydrolysis resulting in a compound of formula (Ia); followed by
- b) cleaving the compound of formula (Ia) of step a under activating conditions in the presence of light resulting in a coumarin (IIa)

In a preferred embodiment, at least the coumarin (IIa) and one of the alcohols (III, IV) are olfactory compounds. Even more preferred are compounds according to the present invention capable of releasing two coumarins (IIa, IIb), herein referred to as Type-II compounds, i.e. compounds of formula (I) wherein n is 1 and R is a monovalent residue of the formula (i) as shown in Scheme 2. Thus Type-II compounds of formula (I) can yield under activating conditions up to four different olfactory compounds.

Scheme 2:



- The activating conditions which lead to the first cleavage step comprise the presence of relative humidity above 20%, preferably above 30% and preferably the presence of a hydrolase such as lipase, esterase, protease or cytochrome P450.


The activating conditions which lead to the second cleavage step comprise the presence of light having a wavelength range of 200 nm to 800 nm, although irradiation with light having in its spectrum wavelengths from 250 nm to 400 nm is preferred. The release of the coumarin of formula IIa/IIb and an alcohol YOH/XOH, or for Y/X = -  
 5 CR<sup>8</sup>=CR<sup>9</sup>R<sup>10</sup> a ketone or aldehyde by tautomerisation, occurs, for example, upon exposure to sunlight penetrating through ordinary windows. Needless to say, it is upon exposure to bright sunlight, especially outdoors, that the release of these compounds will occur faster and to a greater extent than upon exposure to interior light of natural or artificial origin. The cleavage of the compound of formula Ia or Ib can also be initiated  
 10 by an appropriate artificial light source, for example a sun-tanning lamp.

The compounds of formula (I) are virtually odourless and insoluble in water, i.e. the water solubility is equal to or smaller than 10 ppm.

15 It has been found that the use of compounds of formula (I) solves the discoloration problem. In addition, the use of compounds of the present invention also results in a very high deposition rate. Especially good results are obtainable by using Type-II compounds of the present invention. Type-II compounds of the present invention are preferred wherein R<sup>4</sup>=R<sup>11</sup>, R<sup>3</sup>=R<sup>13</sup> and R<sup>2</sup>=R<sup>12</sup>. These molecules deposit on a fabric up  
 20 to 100% by weight based on the amount added to the rinse cycle.

Most of the compounds of the present invention have never been described in literature and thus are novel in its own right.

25 Accordingly the present invention refers in a further aspect to a compound of formula (I)



wherein the acrylic acid ester double bond is of the E configuration;

n is zero or 1;


- 5     Y is  $-\text{CR}^5\text{R}^6\text{R}^7$ , wherein  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^7$  are independently hydrogen or a  $\text{C}_1$ -  $\text{C}_{18}$ , preferably  $\text{C}_1$ -  $\text{C}_{10}$ , hydrocarbon residue of which preferably at least one residue  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^7$  is not hydrogen, and the sum of all carbon atoms ( $\text{R}^5 + \text{R}^6 + \text{R}^7$ ) is not greater than 18 and at least 6, preferably the sum of all carbon atoms is between 6 and 15; or
- 10   Y is  $-\text{CR}^5\text{R}^6\text{R}^7$ , wherein  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^7$  are independently hydrogen or a  $\text{C}_1$ -  $\text{C}_{18}$ , preferably  $\text{C}_1$ -  $\text{C}_{10}$ , aliphatic residue containing one or more atoms/groups selected from O, N and C(O), of which preferably at least one residue  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^7$  is not hydrogen, and the sum of all carbon atoms ( $\text{R}^5 + \text{R}^6 + \text{R}^7$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15, for example Y is 2-(2-butoxy-ethoxy)-ethyl; or
- 15   Y is  $-\text{CR}^8=\text{CR}^9\text{R}^{10}$ , wherein  $\text{R}^8$ ,  $\text{R}^9$  and  $\text{R}^{10}$  are independently hydrogen or a  $\text{C}_1$ -  $\text{C}_{18}$ , preferably  $\text{C}_1$ -  $\text{C}_{10}$ , hydrocarbon residue, of which preferably at least one of the residues  $\text{R}^8$ ,  $\text{R}^9$  and  $\text{R}^{10}$  is not hydrogen, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $\text{R}^8 + \text{R}^9 + \text{R}^{10}$ ) is not greater than 18, preferable the sum of all
- 20   carbon atoms is between 6 and 15; or
- Y is  $-\text{CR}^8=\text{CR}^9\text{R}^{10}$ , wherein  $\text{R}^8$ ,  $\text{R}^9$  and  $\text{R}^{10}$  are independently hydrogen or a  $\text{C}_1$ -  $\text{C}_{18}$ , preferably  $\text{C}_1$ -  $\text{C}_{10}$ , hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), of which preferably at least one of the residues  $\text{R}^8$ ,  $\text{R}^9$  and  $\text{R}^{10}$  is
- 25   not hydrogen, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $\text{R}^8 + \text{R}^9 + \text{R}^{10}$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15;
- $\text{R}^2$  and  $\text{R}^3$  are independently hydrogen;  $\text{C}_1$ - $\text{C}_6$  alkyl, e.g. methyl, ethyl, iso-propyl, n-butyl, tert-butyl;  $\text{C}_1$ - $\text{C}_6$  alkoxy residue, e.g. methoxy, ethoxy;  $-\text{NO}_2$ ;  $-\text{NH}_2$ ;  $-\text{NHCO}_2\text{CH}_3$ ;  $-\text{N}(\text{C}_1\text{-}\text{C}_6\text{ alkyl})_2$ , e.g. dimethylamino, diethylamino;  $-\text{N}(\text{hydroxyalkyl})_2$ , e.g. di(hydroxyethyl)amino, di(hydroxypropyl)amino;  $-\text{NHC}(\text{O})-(\text{C}_1\text{-}\text{C}_8\text{ alkyl})$  or  $-\text{NHC}(\text{O})-(\text{C}_3\text{-}\text{C}_8\text{ aryl})$ , e.g.  $-\text{NHC}(\text{O})$ -methyl or  $-\text{NHC}(\text{O})$ -phenyl; or
- 30

$R^2$  and  $R^3$  are attached at the positions C(6,7), C(7,8), or C(8,9), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

$R^4$  in 2- or 3-position is hydrogen;  $C_1$ - $C_4$  alkyl, e.g. methyl, ethyl, tert-butyl;  $C_2$ - $C_4$  alkenyl, e.g. vinyl, propenyl;  $C_3$ - $C_6$  cycloalkyl, e.g. cyclopropyl, cyclopentyl, cyclohexyl; or  $-CN$ ; and

if  $n$  is zero,  $R$  is a  $C_2$ - $C_{24}$ , preferably  $C_2$ - $C_{18}$ , hydrocarbon residue, e.g. ethyl or phenyl; or  $C_1$ - $C_{24}$ , preferably  $C_1$ - $C_{18}$ , hydrocarbon residue containing one or more heteroatoms selected from N, O and Si; or

if  $n$  is 1,  $R$  is a  $C_1$ - $C_{25}$ , preferably  $C_1$ - $C_{18}$ , hydrocarbon residue; a  $C_1$ - $C_{25}$  hydrocarbon residue containing one or more atoms/groups selected from N, O, Si, and C(O); or  $C_1$ - $C_{25}$ , preferably  $C_1$ - $C_{18}$ , hydrocarbon residue substituted by an ionic substituent of the formula  $N(R^{20})_3^+$ , in which  $R^{20}$  is the residue of an alkyl group with 1 to 18 carbon atoms, preferably 1 to 8 carbon atoms, such as trimethylammonium, or tributylammonium; or  $R$  is a monovalent residue of the formula (i)



wherein

$X$  is  $-CR^{14}R^{15}R^{16}$ , wherein  $R^{14}$ ,  $R^{15}$  and  $R^{16}$  are independently hydrogen or a  $C_1$ - $C_{18}$ , preferably  $C_1$ - $C_{10}$ , hydrocarbon residue, of which preferably at least one residue  $R^{14}$ ,  $R^{15}$  and  $R^{16}$  is not hydrogen, and the sum of all carbon atoms ( $R^{14} + R^{15} + R^{16}$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15; or

$X$  is  $-CR^{14}R^{15}R^{16}$ , wherein  $R^{14}$ ,  $R^{15}$  and  $R^{16}$  are independently hydrogen or a  $C_1$ - $C_{18}$ , preferably  $C_1$ - $C_{10}$ , hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), of which preferably at least one residue  $R^{14}$ ,  $R^{15}$  and



$R^{16}$  is not hydrogen, and the sum of all carbon atoms ( $R^{14} + R^{15} + R^{16}$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15, for example X is 2-(2-butoxy-ethoxy)-ethyl; or

5 X is  $-CR^{17}=CR^{18}R^{19}$ , wherein  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  are independently hydrogen or a  $C_1$ - $C_{18}$ , preferably  $C_1$ - $C_{10}$ , hydrocarbon residue, of which preferably at least one of the residues  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  is not hydrogen, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^{17} + R^{18} + R^{19}$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15; or

10 X is  $-CR^{17}=CR^{18}R^{19}$ , wherein  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  are independently hydrogen or a  $C_1$ - $C_{18}$ , preferably  $C_1$ - $C_{10}$ , hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), of which preferably at least one of the residues  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  is not hydrogen, the geometry of the enol double bond is E or Z, and  
15 the sum of all carbon atoms ( $R^{17} + R^{18} + R^{19}$ ) is not greater than 18, preferable the sum of all carbon atoms is between 6 and 15;

$R^{12}$  and  $R^{13}$  are independently hydrogen;  $C_1$ - $C_6$  alkyl, e.g. methyl, ethyl, iso-propyl, n-butyl, tert-butyl;  $C_1$ - $C_6$  alkoxy residue, e.g. methoxy, ethoxy;  $-NO_2$ ;  $-NH_2$ ;  
20  $-NHCO_2CH_3$ ;  $-N(C_1-C_6 \text{ alkyl})_2$ , e.g. dimethylamino, diethylamino;  $N(\text{hydroxyalkyl})_2$ , e.g. di(hydroxyethyl)amino, di(hydroxypropyl)amino;  $-NHC(O)-(C_1-C_8 \text{ alkyl})$ ; or  $-NHC(O)-(C_3-C_8 \text{ aryl})$ , e.g.  $-NHC(O)$ -methyl or  $-NHC(O)$ -phenyl; or

$R^{12}$  and  $R^{13}$  are attached at the positions C(vi,vii), C(vii,viii), or C(viii,ix), and form  
25 together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

$R^{11}$  in ii- or iii-position is hydrogen;  $C_1$ - $C_4$  alkyl, e.g. methyl, ethyl, tert-butyl;  $C_2$ - $C_4$  alkenyl, e.g. vinyl, propenyl;  $C_3$ - $C_6$  cycloalkyl, e.g. cyclopropyl, cyclopentyl,  
30 cyclohexyl; or  $-CN$ .

As used in relation to the compounds of formula (Ia) "hydrocarbon residue" unless  
otherwise indicated refers to aliphatic residues, e.g. alkyl, alkenyl, alkynyl, and alicyclic  
residues such as cycloalkyl, cycloalkenyl alkylcycloalkyl, alkenylcycloalkyl,  
35 alkenylcycloalkenyl, aryl, alkylaryl or arylalkyl, and "hydrocarbon residues containing

one or more atoms/groups selected from O, N and C(O)," refers to aliphatic residues, e.g. alkyl, alkenyl, alkynyl, and alicyclic residues such as cycloalkyl, cycloalkenyl alkylcycloalkyl, alkenylcycloalkyl, alkenylcycloalkenyl, aryl, alkylaryl or arylalkyl wherein one or more carbon atoms are replaced by O, N and/or C(O).

5

The compounds of formula (I) are advantageously prepared from the corresponding *E*-3-(2-Hydroxy-phenyl)acrylic acid esters, which in turn can be prepared for example via the following methods.

10 In a first step, a corresponding alcohol HO-C(R<sup>4</sup>R<sup>5</sup>R<sup>6</sup>) is transformed into its Li-, Na- or K-salt, preferably its Na-salt, via procedures known to the person skilled in the art, then the corresponding coumarin of formula IIa or IIb is added and the mixture is allowed to react at elevated temperature (20-120°C, preferably 50-100°C) until complete conversion of the coumarin. Standard acid hydrolysis and workup yields the corresponding *E*-3-(2-Hydroxy-phenyl)acrylic acid ester, according to the general  
15 procedure described by Ganguly, N. et al; Synthetic Communications 2001, 31(2), pages 301-309.

Alternatively, salicyl aldehyde is reacted with a dialkoxyposphoryl acetic acid ester of HO-C(R<sup>4</sup>R<sup>5</sup>R<sup>6</sup>), prepared via the Arbuzov reaction between the corresponding chloro- or bromoacetic acid esters and a phosphoric acid trialkyl ester, under the conditions of a  
20 Horner-reaction known to the person skilled in the art.

The preparation of coumarins of formula IIa or IIb are well known to the person skilled in the art and is described for example by A.G. Osborne et al., J. Chem. Research (S), 2003, 114 – 115.

25 The resulting *E*-3-(2-hydroxy-phenyl)acrylic acid ester prepared according to one of the methods described above is then acylated in a second step with an activated acid derivative, such as an acid halide or an acid anhydride, or a corresponding chloroformate under standard conditions well known to the person experienced in organic synthesis.

30 The compounds of formula (I) can be used in any product in which a prolonged and defined release of the abovementioned fragrant compounds is desired. Therefore, these compounds are especially useful in functional perfumery, in products which are exposed to (sun) light during or after application.

The compounds of formula (I) can act as fragrance precursors in functional and fine perfumery i.e. in fine fragrances, industrial, institutional, home and personal care products. Industrial, institutional and home cleaning products to which the compound of formula (I) can be added include all kinds of detergents, window cleaners, hard surface  
5 cleaners, all-purpose cleaners and furniture polishes. Preferably, the products are liquids, e.g. fabric conditioner compositions. A substrate, such as a fabric, treated with a product comprising a compound of formula (I) will diffuse a fresh and/or clean odor under cleavage conditions for much longer than when treated with a conventional product. Fabrics or clothes washed with such fabric softener will release the coumarins  
10 and alcohols, aldehydes or ketones even after having been stored for weeks in a dark place, e.g. a wardrobe.

The compounds of the formula (I) are also useful for application in all kinds of body care products. Especially interesting products are hair care products, for example shampoos,  
15 conditioners and hairsprays, and skin care products such as cosmetic products and especially sun protection products.

The abovementioned examples are of course only illustrative and non-limiting. Many other products to which the compounds of formula (I) may be added include soaps,  
20 bath and shower gels, deodorants and even perfumes and colognes.

The compounds of formula (I) can be used alone or in combination with other fragrance ingredients, solvents or adjuvants known to those skilled in the art. Such ingredients are described, for example, in "Perfume and Flavor Chemicals", S. Arctander, Ed., Vol. I &  
25 II, Allured Publishing Corporation, Carol Stream, USA, 2003 and include fragrance compounds of natural or synthetic origin and essential oils.

The amounts in which the compounds of formula (I) are incorporated in the various above-mentioned products vary within a wide range. The amounts depend on the  
30 nature of the coumarines and alcohols to be released, the nature of the product to which the compounds of formula (I) are added and the desired olfactory effect. The amounts used also depend on the co-ingredients in a given composition when the compounds of formula (I) are used in admixture with perfuming co-ingredients, solvents or adjuvants. Typical concentrations are in the order of 0.01% to 5% by weight of the  
35 products.

The following non-limiting examples further illustrate the embodiments of the invention.

5    Example 1: Preparation of 3-(2-Hydroxy-phenyl)acrylic acid 3,7-dimethyl-oct-6-enyl ester

To a suspension of NaH (114 g of a 60%-dispersion in mineral oil, 2.85 mol) in toluene (500 ml) is added at room temperature a solution of citronellol (468 g, 3.0 mol) in  
10    toluene (800 ml) over 50 min via dropping funnel. The temperature is raised to 85°C (bath) over 60 min and stirring continued for further 45 min. Then a solution of coumarin (219 g, 1.5 mol) in toluene (800 ml) is added over 75 min. After further 90 min stirring at 80°C (inside temperature), the deep orange suspension is cooled to 55°C and poured on a mixture of 2.5 kg crushed ice and 280 ml 37% aq. HCl-solution. The reaction flask  
15    is rinsed twice with 500 ml toluene. Upon stirring for 10 min the colour of the organic layer fades to a pale yellow. The organic layer is washed twice with water, then with brine/water 1:1 and dried over MgSO<sub>4</sub>. After concentrating in the rotary evaporator the excess citronellol is distilled off using a short path apparatus (110-125°C/0.03 mbar, head 82°C) to obtain 240 g of citronellol and 463 g of a brownish residue. The latter is  
20    dissolved in 600 ml hexane containing 18 ml acetone and crystallized at -25°C. After filtration and drying 320 g (71%) of product are obtained as white crystals, m.p. 37-39°C.

IR (film): 3500-3100br, 1673vs, 1633w, 1619w, 1598s, 1450s.

25    <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) : 8.10 (d, J=16, 2H), 7.79 (s, 1H), 7.44 (dd, J=7.6, 1.2, 1H), 7.22-7.18 (m, 1H), 6.69 (d, J=16, 2H), 5.09 (sym. m, 1H), 4.28 (sym. m, 24H), 2.00 (sym. m, 2H), 1.80-1.15 (series of m, 5H), 1.68 (d, J=0.4, 3H), 1.60 (s, 3H), 0.95 (d, J=6.8, 3H).

<sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) : 169.1 (s), 155.9 (s), 141.3 (d), 131.4 (d), 131.2 (s), 129.1  
30    (d), 124.5 (d), 121.5 (s), 120.3 (d), 117.8 (d), 116.4 (d), 63.4 (t), 36.9 (t), 35.3 (t), 29.4 (d), 25.6 (q), 25.3 (t), 19.3 (q), 17.6 (q).

MS (EI 70 eV) : 302 (<1, M<sup>+</sup>), 165 (15), 147 (83), 138 (45), 81 (100).

35    Example 2: Preparation of 3-(2-Hydroxy-phenyl)acrylic acid dec-9-enyl ester

To a suspension of NaH (271 g of a 60%-dispersion in mineral oil, 6.81 mol) in toluene (1200 ml) is added at room temperature a solution of 9-decen-1-ol (1060 g, 6.81 mol) in toluene (1500 ml) over 75 min via dropping funnel. After 10 min stirring a solution of coumarin (495 g, 3.39 mol) in toluene (1800 ml) is added over 75 min. The temperature is raised to 85°C (bath) over 45 min. After further 90 min stirring at 80°C (inside temperature), the deep orange suspension is cooled to 50°C and poured on a mixture of 4 l 10% aq. H<sub>2</sub>SO<sub>4</sub>-solution and 2 l of MTBE. The organic layer is washed with saturated aq. NaHCO<sub>3</sub>-solution, followed by water and brine. After concentrating i. RV the excess of 9-decen-1-ol is removed by thin film evaporation (130°C, 0.05 mbar) to leave 1064 g of residue which is crystallized from hexane at 5°C. From this, 775 g (75%) of product are obtained as pale yellow crystals, m.p. 58°C.

IR (film): 3197br, 1670vs, 1598vs, 1451s.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) : 8.09 (d, J=16, 2H), 7.64 (s, 1H), 7.45 (d, J=8, 1H), 7.23-7.19 (m, 1H), 6.90-6.86 (m, 2H), 6.68 (d, J=16, 1H), 5.84-5.77 (sym. m, 1H), 5.01-4.91 (m, 2H), 4.23 (t, J=8, 2H), 2.03 ("q", J=8, 2H), 1.71 (quint, J=8, 8), 1.39-1.29 (m, 1H).

<sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) : 169.0 (s), 155.8 (s), 141.1 (d), 139.1 (d), 131.4 (d), 129.1 (d), 121.5 (s), 120.3 (d), 117.8 (d), 116.4 (d), 114.0 (t), 65.0 (t), 33.7 (t), 29.3 (t), 29.3 (t), 29.1 (t), 29.0 (t), 28.8 (t), 28.6 (t), 25.8 (t).

MS (EI 70 eV) : 302 (10, M<sup>+</sup>), 164 (8), 178 (100), 118 (37).

Example 3: 3-{2-[2-(2-Dec-9-enyloxycarbonyl-vinyl)phenoxy]carbonyloxy}phenyl}acrylic acid dec-9-enyl ester

25

To a suspension of NaH (14.9 g of a 60% dispersion in mineral oil, 0.31 mol; 1.1 equiv., oil washed away with hexane under argon atmosphere) in toluene (93 ml) is added at room temperature the solution of 9-decen-1-ol (53.2 g, 0.341 mol, 1.1 equiv.) in toluene (93 ml). The solution of coumarin (45.3 g, 0.31 mol, 1.0 equiv.) in toluene (93 ml) is then added within 30 min via dropping funnel. The temperature of the oilbath is raised slowly to 85°C within 30 min, upon which steady evolution of hydrogen is observed. During the following 3 h stirring at 85°C a gelatine-like orange mixture is formed which is then cooled to room temperature. The solution of phosgene in toluene (20%, 100 ml, 0.18 mol 0.6 equiv.) is added over 45 min. During the addition, the gel becomes liquid again and the reaction is cooled with an icebath. The mixture is left stirring for 16 h at

35

room temperature, then the excess phosgene is removed by purging with Argon. The mixture is poured on 200 ml 2N aq. HCl and 200 g ice. The phases are separated and the organic layer is washed three times each with water and water/brine 1:1. After drying the over  $\text{MgSO}_4$ , the volatiles are removed i. RV and the residue dried at 0.05 mbar/50°C for 30 min. From this, 107.1 g of product were obtained as a pale yellow oil which contains the 83% of the title compound (= 90% yield) besides some mixed carbonates. A sample is further purified via column flash chromatography on  $\text{SiO}_2$  eluting with hexane/MTBE 4:1 to isolate analytically pure product as a colourless oil.

IR (film): 1784m, 1713s, 1638m, 1202vs, 758s.

$^1\text{H-NMR}$  (400 MHz,  $\text{CDCl}_3$ ): 7.89 (d,  $J=16$ , 2H), 7.67 (dd,  $J=7.6$ , 1.2, 2H), 7.46-7.30 (m, 6H), 6.53 (d,  $J=16$ , 2H), 5.08 (sym. m, 2H), 4.26 (sym. m, 4H), 1.99 (sym. m, 4H), 1.80-1.15 (4 series of m, 10H), 1.67 (s, 6H), 1.59 (s, 6H), 0.95 (d,  $J=6.4$ , 6H).

$^{13}\text{C-NMR}$  (100 MHz,  $\text{CDCl}_3$ ): 166.5 (s), 151.3 (s), 149.2 (s), 137.3 (d), 131.3 (s), 131.2 (d), 128.1 (d), 127.0 (s), 126.9 (d), 124.5 (d), 122.3 (d), 121.3 (d), 63.3 (t), 37.0 (t), 35.5 (t), 29.6 (d), 25.7 (q), 25.4 (t), 19.4 (q), 17.6 (q).

MS (EI 70 eV): 631 (5,  $\text{M}^+$ ), 493 (10), 475 (100), 431 (26), 337 (65).

**Example 4: 3-(2-Dec-9-enyloxycarbonyloxy-phenyl)acrylic acid dec-9-enyl ester**

**a) 3-(2-Chlorocarbonyloxy-phenyl)acrylic acid dec-9-enyl ester**

The solution of 3-(2-Hydroxy-phenyl)-acrylic acid dec-9-enyl ester (prepared in Example 2, 47.1 g, 156 mmol) in toluene (550 ml) is cooled with an ice bath, then phosgene is introduced via dropping funnel (100 ml of a 20% solution, 188 mmol, 1.2 equiv), followed by  $\text{N,N}$ -diethylanilin (27.5 ml, 169 mmol). The resulting turbid solution is stirred at room temperature for 16 h, after which the excess phosgene is removed by extensive purging with argon. The white suspension is poured on a mixture of 2 N aq. HCl-solution and ice, the phases are separated and the aqueous layer is extracted further with toluene. The organic layers are washed with 2 N aq. HCl-solution, then 3 times with brine and combined. After drying over  $\text{MgSO}_4$ , the solvent is removed and the residue dried at 0.05 mbar/50°C for 20 min. From this 57.2 g (100%) of product are obtained as a pale yellow oil which is not further purified.

30084 PCT /07.02.05



IR (film): 1786s, 1714s, 1639m, 1171s, 1108vs.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>): 7.77 (d, J=16, 1 H), 7.66 (d, J=8, 1 H), 7.44 (m, 1 H), 7.35  
5 (t, J=7, 1 H), 7.27 (d, J=8, 1 H), 6.50 (d, J=16, 1 H), 5.81 (m, 1 H), 4.99 (dd, J=17, 1, 1  
H), 4.93 (dd, J=10, 1, 1 H), 4.22 (t, J=7, 2 H), 2.04 (q, J=7, 2 H), 1.71 (qd, J=7, 7, 2 H),  
1.36 (br. m, 10 H).

<sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>): 166.2 (s), 149.6 (s), 149.2 (s), 139.0 (d), 136.4 (d), 131.2  
10 (d), 128.1 (d), 127.6 (d), 126.6 (s), 121.8 (d), 121.8 (d), 114.1 (t), 64.9 (t), 33.7 (t), 29.3  
(t), 29.1 (t), 28.9 (t), 28.8 (t), 28.6 (t), 25.8 (t).

MS (EI 70 eV): 344 (4, M<sup>+</sup>), 227 (6), 209 (50), 181 (15), 147 (100), 118 (29).

b) 3-(2-Dec-9-enyloxycarbonyloxy-phenyl)acrylic acid dec-9-enyl ester

15

The chloroformate prepared above (3.65 g, 10 mmol) in toluene (10 ml) is added  
dropwise to the solution of 9-decen-1-ol (1.56 g, 10 mmol) and pyridine (2.0 ml, 25  
mmol) in toluene (20 ml) with external ice cooling. The resulting white suspension is  
warmed to room temperature and stirred for 4 h, then 2 N aq. HCl-solution (20 ml) is  
20 added and stirring continued for 5 min. The mixture is then transferred into a separatory  
funnel, the phases are separated and the aqueous phase is further extracted with  
MTBE. The organic layers are washed with brine/H<sub>2</sub>O 1:1 and dried over MgSO<sub>4</sub>.  
Removal of the solvents and drying at 0.05 mbar (at room temperature) yields a crude  
which is purified via flash chromatography on SiO<sub>2</sub> eluting with MTBE/hexane 4:1. From  
25 this, 3.89 g (80%) of product are obtained as a colourless viscous oil.

IR (film): 2926m, 1764s, 1715s, 1639w, 1212vs.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) : 7.82 (d, J=16, 1H), 7.62 (dd, J=7.6/1.2, 1H), 7.40 (br. t,  
J=8, 1H), 7.29-7.20 (m, 2H), 6.48 (d, J=16, 1H), 5.85-5.78 (m, 2H), 5.02-4.92 (m, 4H),  
30 4.26 (t, 6.8, 2H), 4.19 (t, J=6.8, 2H), 2.07-2.02 (m, 4H), 1.80-1.68 (m, 4H), 1.38-1.32 (m,  
20H).

<sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) : 166.6 (s), 153.2 (s), 149.4 (s), 139.0 (d), 139.0 (d), 137.6  
(d), 130.9 (d), 127.7 (d), 127.03 (s), 126.3 (d), 122.5 (d), 120.6 (d), 114.0 (t), 114.0 (t),  
69.2 (t), 64.6 (t), 33.6 (2 t), 29.2 (t), 29.2 (t), 29.1 (t), 29.0 (t), 29.8 (t), 28.9 (t), 28.8 (t),  
35 28.6 (t), 28.4 (t), 25.8 (t), 25.5 (t).

MS (EI 70 eV) : 484 (3, M<sup>+</sup>), 328 (3), 302 (3), 164 (14), 146 (100), 118 (20).



Example 5: 3-[2-(3,7-Dimethyl-oct-6-enyloxycarbonyloxy)phenyl]acrylic acid 3,7-dimethyl-oct-6-enyl ester

- 5 The solution of 3-(2-Hydroxy-phenyl)acrylic acid 3,7-dimethyl-oct-6-enyl ester (6.04 g 20 mmol, as prepared in Example 1) and pyridine (4.0 ml, 50 mmol) in toluene (40 ml) is cooled with an icebath and the solution of citronellyl chloroformate (5.06 g, 22 mmol) in toluene (20 ml) is added dropwise via dropping funnel over 20 min. The resulting suspension is stirred at room temperature during 3 days, then quenched by addition of
- 10 2 N aq. HCl-solution at 2-3°C. Extraction, separation of phases, washing of the organic layer with brine and drying over MgSO<sub>4</sub>, followed by removal of the solvents and drying of the residue under high vacuum yielded 9.7 g (100%) of analytically pure product as a pale yellow oil.
- 15 IR (film): 1765m, 1715m, 1638w, 1213vs.  
<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) : 7.81 (d, J=16, 1H), 7.62 (dd, J=1.2 and 7.6, 1H), 7.29-7.20 (m, 2H), 6.47 (d, J=16, 1H), 5.10 (sym. m, 2H), 4.32-4.22 (m, 4H), 2.00 (sym. m, 4H), 1.85-1.15 (series of m, 10H), 1.68 ("s", 6H), 1.59 (s, 6H), 0.96 (d, J=6, 3H), 0.94 (d, J=6, 3H).
- 20 <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) : 166.5 (s), 153.2 (s), 149.4 (s), 137.6 (d), 131.3 (s), 131.2 (s), 130.9 (d), 127.8 (d), 127.0 (s), 126.3 (d), 124.4 (d), 124.3 (d), 122.5 (d), 120.6 (d), 67.7 (t), 63.1 (t), 36.9 (t), 36.8 (t), 35.4 (t), 35.2 (t), 29.5 (d), 29.2 (d), 25.6 (q), 25.3 (t), 25.2 (t), 19.3 (q), 19.2 (q), 17.5 (q).  
MS (APCI with NH<sub>4</sub>OAc, pos.) : 502 ([M+NH<sub>4</sub>]<sup>+</sup>), 485 (12, M<sup>+</sup>).
- 25

Example 6: 3-(2-Hex-3-enyloxycarbonyloxy-phenyl)acrylic acid 3,7-dimethyl-oct-6-enyl ester

- 30 Repeating the procedure of Example 5 with 3-(2-Hydroxy-phenyl)acrylic acid 3,7-dimethyl-oct-6-enyl ester (6.04 g 20 mmol), pyridine (4.0 ml, 50 mmol) and cis-3-hexenyl chloroformate (3.58 g, 22 mmol) yields 8.60 g (100%) analytically pure product as a pale yellow oil.
- 35 IR (film): 1765m, 1715m, 1638w, 1212vs.

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) : 7.81 (d, J=16, 1H), 7.62 (dd, J=1.2 and 7.6, 1H), 7.39 (sym. m, 1H), 7.29-7.20 (m, 2), 6.47 (d, J=16, 1H), 5.56 (sym. m, 1H), 5.36 (sym. m, 1H), 5.10 (sym. m, 1H), 4.27-4.23 (m, 4H), 2.52 (q, J=8, 4H), 2.11-1.90 (m, 4H), 1.80-1.15 (series of m, 5H), 1.68 (s, 3H), 1.61 (s, 3H), 0.97 (t, J=8, 3H), 0.95 (d, J=8, 3H).  
5 <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) : 166.5 (s), 153.2 (s), 149.4 (s), 137.5 (d), 135.2 (d), 131.2 (s), 130.9 (d), 127.8 (d), 127.0 (s), 126.3 (d), 124.4 (d), 122.5 (d), 122.5 (d), 120.6 (d), 68.4 (t), 63.1 (t), 36.9 (t), 35.4 (t), 29.5 (d), 26.5 (t), 25.6 (q), 25.3 (t), 19.3 (q), 17.5 (q), 14.0 (q).  
MS (APCI with NH<sub>4</sub>OAc, pos.) : 446 ([M+NH<sub>4</sub>]<sup>+</sup>), 429 (11, M<sup>+</sup>).

10

Example 7: 3-(2-Acetoxy-phenyl)acrylic acid dec-9-enyl ester

3-(2-Hydroxy-phenyl)acrylic acid dec-9-enyl ester (6.07 g, 20 mmol, as prepared in  
15 Example 2) in toluene (40 ml) are added to a suspension of NaH (840 mg of a 60% dispersion in mineral oil, 20 mmol, 1 equiv.) in toluene (60 ml). The yellow suspension is stirred for 30 min. at RT, then acetyl chloride (1.96 g, 25 mmol, 1.25 equiv.) are added. The resulting colourless solution is stirred for another 60 min. at RT, then poured on H<sub>2</sub>O (100 ml). The organic layer is separated, and the aqueous layer  
20 extracted with MTBE. The organic layers are washed with 1 N aq. NaHCO<sub>3</sub>-solution, then water and brine. After drying over MgSO<sub>4</sub>, the solvents are removed to yield 6.85 g (99%) of product as a colourless oil.

IR (film): 1770m, 1713s, 1638w, 1170vs.  
25 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ ppm 1.34 (m, 9H), 1.68 (qd, J=7, 7, 2H), 2.03 (q, J=7, 2H), 2.34 (s, 3H), 4.18 (t, J=7, 2H), 4.93 (m, 2H), 5.80 (m, 1H), 6.45 (d, J=16, 1H), 7.10 (m, 1H), 7.22 (t, J=7.58, 1H), 7.36 (m, 1H), 7.62 (dd, J=8, 2, 1H), 7.75 (d, J=16, 1H).  
MS (EI 70 eV): 344 (3, M<sup>+</sup>), 302 (18), 285 (10), 164 (12), 146 (100), 136 (11), 118 (30).

30

Example 8: 3-(2-Hydroxy-phenyl)-acrylic acid phenethyl ester

The compound is prepared by reaction of coumarin with phenylethanol in the presence of sodium hydride following the procedure described in Example 2. The title compound  
35 is isolated as a white solid, m.p. 40°C.

IR (film): 3261 vs, 1771m, 1738s, 1679vs, 1631vs, 1599 vs, 1174 vs, 752 vs, 700 vs.  
<sup>13</sup>C-NMR (CDCl<sub>3</sub>, 100 MHz): 168.6 (s), 155.8 (s), 141.3 (d), 137.7 (d), 131.5 (d), 129.1  
(d), 128.9 (d), 128.5 (d), 126.5 (d), 121.5 (s), 120.4 (d), 117.7 (d), 116.4 (d), 65.2 (t),  
5 35.1 (t).  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz): 8.07 (d, *J*=16, 1 H), 7.49 (s, 1 H), 7.42 (dd, *J*=8, 2, 1 H),  
7.27 - 7.32 (m, 2 H), 7.17 - 7.26 (m, 5 H), 6.86 (td, *J*=8, 1.1, 2 H), 6.64 (d, *J*=16, 1 H),  
4.43 (t, *J*=7, 2 H), 3.01 (t, *J*=7, 2 H).  
MS (EI, 70 eV): 268 (<1, M<sup>+</sup>), 266 (1), 250 (4), 164 (46), 146 (69), 118 (60), 104 (79), 91  
10 (100).

Example 9: 3-(2-Phenethyloxycarbonyloxy-phenyl)-acrylic acid phenethyl ester

15 A solution of 3-(2-Hydroxy-phenyl)-acrylic acid phenethyl ester (1.61 g, 6.0 mmol) in  
toluene (15 ml) is added dropwise at room temperature to a suspension formed from  
chloroformic acid phenethyl ester (1.22 g, 6.6 mmol, 1.1 equiv.) and pyridine (0.97 ml,  
12.0 mmol, 2.0 equiv.) in toluene (10 ml). The mixture is further stirred at room  
temperature for 16 h, then worked up following the general procedure described for  
20 Example 5 and purified by flash chromatography on SiO<sub>2</sub>, eluting with hexane/MTBE  
4:1. The title compound (1.61 g, 64% yield) is obtained as a colourless, viscous oil.

IR (film): 1760s, 1710 s, 1212 vs, 1168 vs, 697 vs.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz): 7.82 (d, *J*=16.2 Hz, 1 H), 7.60 (dd, *J*=7.8, 1.6 Hz, 1 H), 7.36  
25 - 7.43 (m, 1 H), 7.20 - 7.35 (m, 11 H), 7.16 (dd, *J*=8.2, 1.1 Hz, 1 H), 6.46 (d, *J*=16.2 Hz,  
1 H), 4.37 - 4.51 (m, 4 H), 2.97 - 3.10 (m, 4 H).  
<sup>13</sup>C-NMR (CDCl<sub>3</sub>, 100 MHz): 166.5 (s), 153.2 (s), 149.5 (s), 137.9 (s), 137.8 (s), 136.8  
(d), 131.1 (d), 129.7 (d), 129.0 (d), 128.9 (d), 128.9 (d), 128.6 (d), 128.6 (d), 128.5 (d),  
127.8 (d), 127.8 (d), 127.0 (s), 126.8 (d), 126.6 (d), 126.5 (d), 122.6 (d), 120.5 (d), 69.4  
30 (t), 65.1 (t), 35.2 (t), 35.0 (t).  
MS (EI, 70 eV): 417 (<1, [M+1]<sup>+</sup>), 370 (4), 164 (8), 146 (7), 105 (100), 104 (31).

Example 10: 3-[2-(3-Methyl-5-phenyl-pentyloxycarbonyloxy)-phenyl]-acrylic acid phen-  
35 ethyl ester

Following the general procedure described in Example 9 the title compound is prepared, using 3-(2-hydroxy-phenyl)-acrylic acid phenethyl ester (1.61 g, 6.0 mmol), chloroformic acid 3-methyl-5-phenyl-pentyl ester (1.59 g, 6.6 mmol, 1.1 equiv.) and pyridine (0.97 ml, 12.0 mmol, 2.0 equiv.). 3-[2-(3-Methyl-5-phenyl-pentyloxycarbonyloxy)-phenyl]-acrylic acid phenethyl ester (1.65 g, 58% yield) is obtained as colourless, viscous oil.

IR (film): 1760 s, 1712 s, 1251 s, 1211 vs, 1167 vs, 697 vs.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz): 7.82 (d, J=16.2 Hz, 1 H), 7.61 (dd, J=7.8, 1.6 Hz, 1 H), 7.37 - 7.44 (m, 1 H), 7.13 - 7.35 (m, 13 H), 6.46 (d, J=16.2 Hz, 1 H), 4.41 (t, J=7.1 Hz, 2 H), 4.25 - 4.37 (m, 2 H), 3.00 (t, J=7.1 Hz, 2 H), 1.79 - 1.94 (m, 1 H), 1.46 - 1.76 (m, 5 H), 1.02 (d, J=6.3 Hz, 3 H)

<sup>13</sup>C-NMR (CDCl<sub>3</sub>, 100 MHz): 166.5 (s), 153.3 (s), 149.5 (s), 142.5 (s), 137.9 (d), 137.8 (s), 131.1 (d), 128.9 (d), 128.5 (d), 128.3 (d), 128.3 (d), 127.9 (d), 127.0 (s), 126.6 (d), 126.5 (d), 125.7 (d), 122.6 (d), 120.5 (d), 67.7 (t), 65.1 (t), 38.8 (t), 35.3 (t), 35.2 (t), 33.2 (t), 29.4 (d), 19.4 (q).

MS (EI, 70 eV): 472 (<1, M<sup>+</sup>), 426 (4), 313 (5), 164 (34), 160 (39), 146 (32), 105 (91), 91 (100).

#### Example 11 - 15

Further compounds as listed in Table 1 were prepared according to the general procedure described in the specification.

Table 1

IR (film)	MS (EI 70eV)	NMR ( <sup>1</sup> H NMR (400 MHz, CDCl <sub>3</sub> ) δ in ppm)
Benzoic acid 4-benzoylamino-2-[2-(3,7-dimethyl-oct-6-enyloxycarbonyl)vinyl]phenyl ester (11)		
1713s,	527 ([M+2] <sup>+</sup> ),	0.87 (m, 3H), 1.15 (m, 1H), 1.31 (m, 2H), 1.44 (m, 2H), 1.52
1654s,	388 (26), 371	(m, 2H), 1.59 (s, 3H), 1.66 (s, 3H), 1.94 (m, 2H), 4.17 (m,
1525s,	(24), 265 (33),	2H), 5.08 (m, 1H), 6.52 (d, J=16, 1H), 7.25 (d, J=9, 1H),
1196vs,	105 (100).	7.55 (m, 5H), 7.66 (m, 2H), 7.80, (d, J=16, 1H), 7.90 (m, 2
704vs.		H), 8.13 (d, J=3, 1H), 8.23 (dd, J=8, 1, 2H).

3-(2-Acetoxy-5-tert-butyl-phenyl)acrylic acid 3,7-dimethyl-oct-6-enyl ester (12)		
1768m, 1713s, 1182vs.	400 (2, M <sup>+</sup> ), 262 (29, 220 (28), 202 (66), 187 (100), 159 (30).	0.94 (d, J=7, 3H), 1.18-1.80 (series of m, 5H), 1.33 (s, 9H), 1.61 (s, 3H), 1.68 (s, 3H), 2.0 (m, 2H), 2.36 (s, 3H) 4.25 (m, 2H), 5.10 (m, 1H), 6.45 (d, J=16, 1H), 7.04 (d, J=9, 1H), 7.42 (dd, J=9, 2, 1H), 7.62 (d, J=2, 1H), 7.74 (d, J=16, 1H).
3-(2-Acetoxy-phenyl)acrylic acid 3,7-dimethyl-oct-6-enyl ester (13)		
1770m, 1713s, 1637w, 1174vs.	344 (<1, M <sup>+</sup> ), 147 (92), 138 (68), 95 (76), 81 (100).	0.95 (d, J=7, 3H), 1.23 (ddd, J=9, 7, 6, 1H), 1.37 (m, 1H), 1.51 (m, 1H), 1.61 (s, 3H), 1.58-1.80 (m, 2H), 1.68 (s, 3H), 2.01 (m, 2H), 2.37 (s, 3H), 4.24 (m, 2H), 5.10 (ddd, J=7, 6, 1, 1H), 6.45 (d, J=16, 1H), 7.12 (dd, J=8, 1, 1H), 7.26 (m, 1H), 7.40 (td, J=8, 2, 1H), 7.64 (dd, J=8, 2, 1H), 7.74 (d, J=16, 1H).
3-(2-Ethoxycarbonyloxy-phenyl)acrylic acid dec-9-enyl ester (14)		
1763s, 1714s, 1638m, 1249s, 1212vs, 1170vs	374 (4, M <sup>+</sup> ), 285 (4), 192 (3), 164 (7), 146 (100), 118 (34).	1.37 (m, 13H), 1.69 (qd, J=7, 7, 2H), 2.04 (q, J=7, 2H), 4.19 (t, J=7, 2H), 4.33 (q, J=7, 2H), 4.93 (d, J=10, 1H), 4.99 (dd, J=17, 2, 1H), 5.81 (m, 1H), 6.48 (d, J=16, 1H), 7.25 (m, 2H), 7.40 (m, 1H), 7.63 (d, J=8, 1H), 7.82 (d, J=16, 1H).
3-(2-{2-[2-(3,7-Dimethyl-oct-6-enyloxycarbonyl)vinyl]phenoxy}carbonyloxy}phenyl)- acrylic acid 3,7- dimethyl-oct-6-enyl ester (15)		
1784w, 1712s, 1638w, 1165vs.	630 (4, M <sup>+</sup> ), 493 (12), 475 (100), 431 (15), 337 (65), 249 (60), 147 (60).	0.95 (d, J=6, 6 H), 1.18-1.80 (series of m, 10 H), 1.57 (s, 6 H), 1.67 (d, J=1, 6 H), 1.99 (m, 4 H), 4.26 (m, 4 H), 5.08 (m, 2 H), 6.53 (d, J=16, 2 H), 7.31 (m, 4 H), 7.44 (m, 2 H), 7.65 (dd, J=8, 2, 2 H), 7.90 (d, J=16, 2 H).

Example 16: 3-(2-Hydroxy-4-methoxy-phenyl)-acrylic acid 3,7-dimethyl-oct-6-enyl ester

- 5 The compound is prepared by reaction of 7-methoxycoumarin with citronellol in the presence of sodium hydride following the procedure described in Example 2. The title compound is isolated as a viscous, colourless oil.

IR (film): 3331 br., 1705 s, 1676 s, 1611 vs, 1518 m, 1169 vs, 836 w, 802 w.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz): 7.98 (d, *J*=16.2 Hz, 1 H), 7.30-7.39 (m, 2 H), 6.34 - 6.63 (m, 3 H), 5.03 - 5.20 (m, 1 H), 4.18 - 4.37 (m, 2 H), 3.79 (s, 3 H), 2.06 (m, 2 H), 1.10 - 1.87 (m, 5 H), 1.68 (s, 3H), 1.61 (s, 3H), 0.95 (d, *J*=6.6 Hz, 3 H).

MS (EI, 70 eV): 348 (<1, M<sup>+</sup>), 194 (81), 177 (87), 176 (100), 148 (54), 133 (39).

Example 17: 3-(2-Hydroxy-5-methyl-phenyl)-acrylic acid 2-ethyl-4-(2,2,3-trimethyl-cyclopent-3-enyl)-but-2-enyl ester

The compound is prepared by reaction of 6-methylcoumarin with 2-ethyl-4-(2,2,3-trimethyl-cyclopent-3-enyl)-but-2-en-1-ol in the presence of sodium hydride following the procedure described in Example 2. The compound is isolated as a viscous, slightly yellow oil.

IR (film): 3313 s, 3035m, 2958 vs, 1688 s, 1627 s, 1508 m, 1463 s, 1155 s, 816 m, 799 m.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 200 MHz): 8.00 (d, *J*=16.2 Hz, 1 H), 7.17 - 7.36 (m, 1 H), 6.92 - 7.13 (m, 2 H), 6.74 (d, *J*=8.2 Hz, 1 H), 6.61 (d, *J*=16.2 Hz, 1 H), 5.31 - 5.63 (m, 2 H), 5.22 (s, 2 H), 4.66 (s, 1 H), 4.07 (s, 2 H), 2.26 (s, 3H), 1.67 - 2.45 (m, 4 H), 1.62-1.56 (m, 3 H), 0.90 (t, *J*=7.6 Hz, 3H), 0.89 (s, 3 H), 0.78 (s, 3 H).

MS (EI, 70 eV): 368 (<1, M<sup>+</sup>), 190 (17), 161 (53), 108 (100).

Example 18: 3-(2-Hydroxy-4-methoxy-phenyl)-acrylic acid 3-methyl-5-phenyl-pentyl ester

The compound is prepared by reaction of 7-methoxycoumarin with citronellol in the presence of sodium hydride following the procedure described in Example 2. The compound is isolated as a white solid, m.p. 70-73°C.

IR (film): 1760 s, 1712 s, 1251 s, 1211 vs, 1167 vs, 697 vs.


$^1\text{H-NMR}$  ( $\text{CDCl}_3$ , 400 MHz): 8.00 (d,  $J=15.9$  Hz, 1 H), 7.93 (br. s, 1 H), 7.11 - 7.39 (m, 6 H), 6.55 (d,  $J=15.9$  Hz, 1 H), 6.42 - 6.49 (m, 2 H), 4.19 - 4.34 (m, 2 H), 3.74 (s, 3 H), 2.52 - 2.74 (m, 2 H), 1.41 - 1.89 (m, 5 H), 1.00 (d,  $J=6.3$  Hz, 3 H).

$^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ , 100 MHz): 169.4 (s), 162.5 (s), 157.5 (s), 142.6 (s), 141.1 (s), 130.5 (s), 128.3 (d), 125.6 (d), 115.0 (s), 114.9 (d), 106.8 (d), 101.7 (d), 63.0 (t), 55.3 (q), 38.7 (t), 35.4 (t), 33.2 (t), 29.5 (d), 19.4 (q).


MS (EI, 70 eV): 354 (30,  $\text{M}^+$ ), 194 (78), 176 (100), 148 (46), 133 (17), 104 (14), 91 (40).

#### 10 Examples 19:


The compounds of Example 16, 17 and 18 may be used as intermediates for the preparation of further compounds A, B and C according to the present invention. Their preparation may be carried with the corresponding chloroformates following the general procedure of Example 9.



3-[2-(3,7-Dimethyloct-6-enyloxycarbonyloxy)-4-methoxy-phenyl]-acrylic acid 3,7-dimethyloct-6-enyl ester



3-(2-Benzyloxycarbonyloxy-5-methyl-phenyl)-acrylic acid 2-ethyl-4-(2,2,3-trimethyl-cyclopent-3-enyl)-but-2-enyl ester



3-[2-(4-tert-Butylcyclohexyloxycarbonyloxy)-4-methoxy-phenyl]-acrylic acid 3-methyl-5-phenyl-pentyl ester



**Example 20:** UV-Spectra Comparison (protected/non protected fragrance in presence of a fabric conditioner)

Solutions A to D, 5ml each, were prepared using CH<sub>3</sub>CN/H<sub>2</sub>O (3:2) as a solvent and their colour visually judged.

Solution	Fragrance Precursor (each added at 0.1 mg/ml)	Additive (amount)	Colour
A	(E)-3-(2-Hydroxy-phenyl)acrylic acid dec-9-enyl ester (Ex. 2)	none	colourless
B	As above (A)	Fabric conditioner <sup>1)</sup> (30 mg)	Bright yellow
C	3-{2-[2-(2-Dec-9-enyloxycarbonyl- vinyl)phenoxy]phenoxy}- acrylic acid dec-9-enyl ester (Ex. 3)	none	colourless
D	As above (C)	Fabric conditioner <sup>1)</sup> (30 mg)	colourless

<sup>1)</sup> Aqueous fabric conditioner emulsion containing 12.7% wt/wt of active cationic surfactant.

After adding a fabric conditioner to a solution comprising an unprotected fragrance precursor (solution A / B) the solution turned bright yellow, whereas a solution comprising a protected fragrance precursor (solution C / D) remains colourless.

The UV-spectra of solutions A to D were recorded and are depicted in Figure 1. They show that upon addition of the fabric conditioner the highest  $\lambda_{\max}$  at 326 nm (UV) shifts to 394 nm (visible light), whereas the highest  $\lambda_{\max}$  of the protected fragrance precursor remains unchanged at 270 nm upon addition of the fabric conditioner.

**Example 21:** Application in fabric conditioner.

The deposition and cleavage of the compounds of formula (I) on cotton (white towels) in a typical wash/rinse cycle is determined as described in the following. All handling of samples containing a compound of formula (I) is done with as little exposure to light as possible.

An aqueous fabric conditioner emulsion containing 12.7% wt/wt of active cationic surfactant and 0.5 % wt/wt of a compound of formula (I) is prepared. Whereas samples with free o-coumarates turn yellow, samples with protected o-coumarates remain white. A wash/rinse cycle (40°C program) in a standard washing machine is performed with a  
5 1 kg wash load consisting of 25 cotton terry towels, adding the following:

- 1) 62 g of a standard concentrated unperfumed washing powder containing protease, cellulase, and lipase for the washing cycle
- 2) 16 g of fabric conditioner as prepared above for the rinse cycle (containing  
10 80 mg of a compound of formula (I) as indicated in Table 2, compound Ex. No 6, 5, 3, 4 or 14, or a compound of formula (A), i.e. comp. Ex. No. 1 or 2, as comparison as indicated in Table 2.

The towels are removed from the washing machine and dried in the dark for 24 h at  
15 room temperature and 40% relative humidity. Three towels (representing each 4% of the total wash load) are removed and placed individually in a Soxhlet apparatus containing 0.5 l of methylene chloride and extracted for 5 h. The solvent is removed carefully in a rotary evaporator and the residues are standardized to 10 ml acetonitrile solution. These solutions are analyzed by RP-HPLC using a H<sub>2</sub>O/acetonitrile gradient  
20 and UV-detection at 258 nm. The concentrations of "protected" and "unprotected" fragrance precursor per sample are determined via external calibration and a mean value is calculated from the three towels. From this, the mean deposition rate in % of theory compared to the molar amount of protected precursor applied via the fabric conditioner for the two precursor types is calculated. The results are listed in Table 2.

25 Best cleavage rate by lipase is observed for carbonates derived from short chain primary alcohols (Comp. Ex. No. 6) and (Comp. Ex. No 14). The best combined deposition/cleavage result is obtained from the symmetrical carbonate (Comp. Ex. No. 3). The final deposition of free precursor using this compound is higher than by  
30 administration of the corresponding "unprotected" 3-(2-Hydroxy-phenyl)-acrylic acid ester (Comp. Ex. No. 2).


As used herein "unprotected fragrance precursor" means a compound comprising a hydroxyl group, i.e. the prior art compound (A), and "protected fragrance precursor"  
35 means a compound of formula (I) according to the present invention.

Table 2

Comp. Ex. N°	Fragrance precursor	Protecting group	Releases Coumarin and	Modified Prec. [Depos. %]	Free prec. [Depos. %]	Cleavage [%]
1	3-(2-Hydroxy-phenyl)acrylic acid 3,7-dimethyl-oct-6-enyl ester	None	Citronellol	-	45	-
2	3-(2-Hydroxy-phenyl)acrylic acid dec-9-enyl ester	None	9-Decen-1-ol	-	46	-
6	3-(2-Hex-3-enyloxycarbonyloxy-phenyl)acrylic acid 3,7-dimethyl-oct-6-enyl ester	cis-3-hex. carbonate	Citronellol & cis-3-hexenol	2	50	97
5	3-[2-(3,7-Dimethyl-oct-6-enyloxycarbonyloxy)-phenyl]acrylic acid 3,7-dimethyl-oct-6-enyl ester	Citronellyl carbonate	Citronellol	69	11	14
3	3-[2-[2-(2-Dec-9-enyloxycarbonyl-vinyl)phenoxy carbonyloxy]phenyl]-acrylic acid dec-9-enyl ester	Sym. Carbonate	9-Decen-1-ol	34	66	66
4	3-(2-Dec-9-enyloxycarbonyloxy-phenyl)acrylic acid dec-9-enyl ester	Rosalva-Carbonate	9-Decen-1-ol	32	34	68
14	3-(2-Ethoxycarbonyloxy-phenyl)acrylic acid dec-9-enyl ester	Ethyl carbonate	9-Decen-1-ol	5	42	91

Claims

1. Use of a compound of formula (I) as precursor for olfactory compounds compound



wherein the acrylic acid ester double bond is of the E configuration;

n is zero or 1;

Y is  $-\text{CR}^5\text{R}^6\text{R}^7$ , wherein  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^7$  are independently hydrogen or a  $\text{C}_1$ -  $\text{C}_{18}$  hydrocarbon residue, and the sum of all carbon atoms ( $\text{R}^5 + \text{R}^6 + \text{R}^7$ ) is not greater than 18; or

Y is  $-\text{CR}^5\text{R}^6\text{R}^7$ , wherein  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^7$  are independently hydrogen or a  $\text{C}_1$ -  $\text{C}_{18}$  hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), and the sum of all carbon atoms ( $\text{R}^5 + \text{R}^6 + \text{R}^7$ ) is not greater than 18; or

Y is  $-\text{CR}^8=\text{CR}^9\text{R}^{10}$ , wherein  $\text{R}^8$ ,  $\text{R}^9$  and  $\text{R}^{10}$  are independently hydrogen or a  $\text{C}_1$ -  $\text{C}_{18}$  hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $\text{R}^8 + \text{R}^9 + \text{R}^{10}$ ) is not greater than 18; or


Y is  $-\text{CR}^8=\text{CR}^9\text{R}^{10}$ , wherein  $\text{R}^8$ ,  $\text{R}^9$  and  $\text{R}^{10}$  are independently hydrogen or a  $\text{C}_1$ -  $\text{C}_{18}$  hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $\text{R}^8 + \text{R}^9 + \text{R}^{10}$ ) is not greater than 18;

$R^2$  and  $R^3$  are independently hydrogen,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy residue,  $-NH_2$ ,  $-NO_2$ ,  $-NHCO_2CH_3$ ,  $-N(C_1-C_6 \text{ alkyl})_2$ ,  $-N(\text{hydroxyalkyl})_2$ ,  $-NHC(O)-(C_1-C_8 \text{ alkyl})$  or  $-NHC(O)-(C_3-C_8 \text{ aryl})$ ; or

$R^2$  and  $R^3$  are attached at the positions C(6,7), C(7,8), or C(8,9), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

$R^4$  in 2- or 3-position is hydrogen,  $C_1$ - $C_4$  alkyl,  $C_2$ - $C_4$  alkenyl,  $C_3$ - $C_6$  cycloalkyl, or  $-CN$ ; and

- a) if  $n$  is zero,  $R$  is a  $C_1$ - $C_{24}$  hydrocarbon residue, or  $C_1$ - $C_{24}$  hydrocarbon residue containing one or more heteroatoms selected from N, O and Si; or
- b) if  $n$  is 1,  $R$  is a  $C_1$ - $C_{25}$  hydrocarbon residue, a  $C_1$ - $C_{25}$  hydrocarbon residue containing one or more atoms/groups selected from N, O, Si, and C(O), or  $C_1$ - $C_{25}$  hydrocarbon residue substituted by an ionic substituent of the formula  $N(R^{20})_3^+$ , in which  $R^{20}$  is the residue of an alkyl group with 1 to 18 carbon atoms; or  $R$  is a monovalent residue of the formula (i)



wherein

$X$  is  $-CR^{14}R^{15}R^{16}$ , wherein  $R^{14}$ ,  $R^{15}$  and  $R^{16}$  are independently hydrogen or a  $C_1$ - $C_{18}$  hydrocarbon residue, and the sum of all carbon atoms ( $R^{14} + R^{15} + R^{16}$ ) is not greater than 18; or

$X$  is  $-CR^{14}R^{15}R^{16}$ , wherein  $R^{14}$ ,  $R^{15}$  and  $R^{16}$  are independently hydrogen or a  $C_1$ - $C_{18}$  hydrocarbon residue containing one or more atoms/groups selected

from O, N and C(O), and the sum of all carbon atoms ( $R^{14} + R^{15} + R^{16}$ ) is not greater than 18; or

X is  $-CR^{17}=CR^{18}R^{19}$ , wherein  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  are independently hydrogen or a  $C_1$ - $C_{18}$  hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^{17} + R^{18} + R^{19}$ ) is not greater than 18; or

X is  $-CR^{17}=CR^{18}R^{19}$ , wherein  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  are independently hydrogen or a  $C_1$ - $C_{18}$  hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^{17} + R^{18} + R^{19}$ ) is not greater than 18;


$R^{12}$  and  $R^{13}$  are independently hydrogen,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy residue,  $-NO_2$ ,  $-NH_2$ ,  $-NHCO_2CH_3$ ,  $-N(C_1-C_6 \text{ alkyl})_2$ ,  $-N(\text{hydroxyalkyl})_2$ ,  $-NHC(O)-(C_1-C_8 \text{ alkyl})$  or  $-NHC(O)-(C_3-C_8 \text{ aryl})$ ; or

$R^{12}$  and  $R^{13}$  are attached at the positions C(vi,vii), C(vii,viii), or C(viii,ix), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

$R^{11}$  in ii- or iii-position is hydrogen,  $C_1$ - $C_4$  alkyl,  $C_2$ - $C_4$  alkenyl,  $C_3$ - $C_6$  cycloalkyl, or  $-CN$ .

2. A consumer product comprising a compound of formula (I) as defined by claim 1.
3. A process for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) as defined by claim 1.
4. A process of providing an olfactory compound to a substrate comprising the steps:
  - a) cleaving a compound of formula (I) as defined by claim 1 by hydrolysis resulting in a compound of formula (Ia); followed by
  - b) cleaving the compound of formula (Ia) of step a under activating conditions in the presence of light resulting in a coumarin (IIa).

5. A compound of formula (I)



wherein the acrylic acid ester double bond is of the E configuration;

$n$  is zero or 1;

$Y$  is  $-CR^5R^6R^7$ , wherein  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen or a  $C_1$ - $C_{18}$  hydrocarbon residue, and the sum of all carbon atoms ( $R^5 + R^6 + R^7$ ) is not greater than 18 and at least 6; or

$Y$  is  $-CR^5R^6R^7$ , wherein  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen or a  $C_1$ - $C_{18}$  aliphatic residue containing one or more atoms/groups selected from O, N and C(O), and the sum of all carbon atoms ( $R^5 + R^6 + R^7$ ) is not greater than 18; or

$Y$  is  $-CR^8=CR^9R^{10}$ , wherein  $R^8$ ,  $R^9$  and  $R^{10}$  are independently hydrogen or a  $C_1$ - $C_{18}$  hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^8 + R^9 + R^{10}$ ) is not greater than 18; or

$Y$  is  $-CR^8=CR^9R^{10}$ , wherein  $R^8$ ,  $R^9$  and  $R^{10}$  are independently hydrogen or a  $C_1$ - $C_{18}$  hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $R^8 + R^9 + R^{10}$ ) is not greater than 18;

$R^2$  and  $R^3$  are independently hydrogen,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy residue,  $-NH_2$ ,  $-NO_2$ ,  $-NHCO_2CH_3$ ,  $-N(C_1-C_6 \text{ alkyl})_2$ ,  $-N(\text{hydroxyalkyl})_2$ ,  $-NHC(O)-(C_1-C_8 \text{ alkyl})$  or




-NHC(O)-(C<sub>3</sub>-C<sub>8</sub> aryl); or

R<sup>2</sup> and R<sup>3</sup> are attached at the positions C(6,7), C(7,8), or C(8,9), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

R<sup>4</sup> in 2- or 3-position is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, or -CN; and

- a) if n is zero, R is a C<sub>2</sub>-C<sub>24</sub> hydrocarbon residue, or C<sub>1</sub>-C<sub>24</sub> hydrocarbon residue containing one or more heteroatoms selected from N, O and Si; or
- b) if n is 1, R is a C<sub>1</sub>-C<sub>25</sub> hydrocarbon residue, a C<sub>1</sub>-C<sub>25</sub> hydrocarbon residue containing one or more atoms/groups selected from N, O, Si, and C(O), or C<sub>1</sub>-C<sub>25</sub> hydrocarbon residue substituted by an ionic substituent of the formula N(R<sup>20</sup>)<sub>3</sub><sup>+</sup>, in which R<sup>20</sup> is the residue of an alkyl group with 1 to 18 carbon atoms; or R is a monovalent residue of the formula (i)



wherein

X is -CR<sup>14</sup>R<sup>15</sup>R<sup>16</sup>, wherein R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> are independently hydrogen or a C<sub>1</sub>-C<sub>18</sub> hydrocarbon residue, and the sum of all carbon atoms (R<sup>14</sup>+R<sup>15</sup>+R<sup>16</sup>) is not greater than 18; or

X is -CR<sup>14</sup>R<sup>15</sup>R<sup>16</sup>, wherein R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> are independently hydrogen or a C<sub>1</sub>-C<sub>18</sub> hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), and the sum of all carbon atoms (R<sup>14</sup>+R<sup>15</sup>+R<sup>16</sup>) is not greater than 18; or


X is  $-\text{CR}^{17}=\text{CR}^{18}\text{R}^{19}$ , wherein  $\text{R}^{17}$ ,  $\text{R}^{18}$  and  $\text{R}^{19}$  are independently hydrogen or a  $\text{C}_1$ - $\text{C}_{18}$  hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $\text{R}^{17} + \text{R}^{18} + \text{R}^{19}$ ) is not greater than 18; or

X is  $-\text{CR}^{17}=\text{CR}^{18}\text{R}^{19}$ , wherein  $\text{R}^{17}$ ,  $\text{R}^{18}$  and  $\text{R}^{19}$  are independently hydrogen or a  $\text{C}_1$ - $\text{C}_{18}$  hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ( $\text{R}^{17} + \text{R}^{18} + \text{R}^{19}$ ) is not greater than 18;

$\text{R}^{12}$  and  $\text{R}^{13}$  are independently hydrogen,  $\text{C}_1$ - $\text{C}_6$  alkyl,  $\text{C}_1$ - $\text{C}_6$  alkoxy residue,  $-\text{NO}_2$ ,  $-\text{NH}_2$ ,  $-\text{NHCO}_2\text{CH}_3$ ,  $-\text{N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$ ,  $-\text{N}(\text{hydroxyalkyl})_2$ ,  $-\text{NHC}(\text{O})-(\text{C}_1\text{-C}_8 \text{ alkyl})$  or  $-\text{NHC}(\text{O})-(\text{C}_3\text{-C}_8 \text{ aryl})$ ; or

$\text{R}^{12}$  and  $\text{R}^{13}$  are attached at the positions C(vi,vii), C(vii,viii), or C(viii,ix), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

$\text{R}^{11}$  in ii- or iii-position is hydrogen,  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_2$ - $\text{C}_4$  alkenyl,  $\text{C}_3$ - $\text{C}_6$  cycloalkyl, or  $-\text{CN}$ .

*Fig. 1-A*

SUBSTITUTE SHEET (RULE 26)







Fig. 1-B

SUBSTITUTE SHEET (RULE 26)




*Fig. 1-C*

SUBSTITUTE SHEET (RULE 26)







*Fig. 1-D*

SUBSTITUTE SHEET (RULE 26)



A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07C69/618 C11D3/50 A61K7/46

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C11D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, CHEM ABS Data, WPI Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	US 2004/248762 A1 (MCGEE THOMAS ET AL) 9 December 2004 (2004-12-09) *compound GR-85-2666/000 of p. 12*	2, 5
X	CAIRNS, NICHOLAS ET AL: "Syntheses of 6-acylcoumarins via highly regioselective Fries rearrangements. Total syntheses of the linear acylated coumarins geijerin and dehydrogeijerin" TETRAHEDRON, 48(36), 7581-90 CODEN: TETRAB; ISSN: 0040-4020, 1992, XP002329069 compounds 6A, B, D-G ----- -/--	5

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

1 June 2005

Date of mailing of the international search report

13/06/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Österle, C

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; YUN, HYUN CHU ET AL: "Generation of graded index profile of poly(methyl methacrylate) by a photochemical reaction" XP002329070 retrieved from STN Database accession no. 2003:721643 abstract & MACROMOLECULAR RESEARCH , 11(4), 236-240 CODEN: MRAECT; ISSN: 1598-5032, 2003, -----	5
X	PATENT ABSTRACTS OF JAPAN vol. 014, no. 552 (M-1056), 7 December 1990 (1990-12-07) - & JP 02 235683 A (RICOH CO LTD), 18 September 1990 (1990-09-18) abstract; compound 1 -----	5
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ICHIMURA, KUNIHIO ET AL: "Role of E/Z photoisomerization of cinnamate side chains attached to polymer backbones in the alignment photoregulation of nematic liquid crystals" XP002329071 retrieved from STN Database accession no. 1996:528115 abstract & JAPANESE JOURNAL OF APPLIED PHYSICS, PART 2: LETTERS , 35(8A), 1992-L995 CODEN: JAPLD8; ISSN: 0021-4922, 1996, -----	5
A	EP 0 936 211 A (GIVAUDAN ROURE S.A) 18 August 1999 (1999-08-18) cited in the application the whole document -----	1-5
A	EP 0 887 338 A (GIVAUDAN-ROURE SA; GIVAUDAN SA) 30 December 1998 (1998-12-30) the whole document -----	1-4

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CH2005/000074

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2004248762 A1	09-12-2004	NONE	
JP 02235683 A	18-09-1990	JP 2832207 B2	09-12-1998
EP 0936211 A	18-08-1999	EP 0936211 A2	18-08-1999
		AU 725999 B2	26-10-2000
		AU 1643099 A	21-10-1999
		BR 9900443 A	02-05-2000
		CN 1227837 A	08-09-1999
		ID 22095 A	02-09-1999
		JP 2000063328 A	29-02-2000
		SG 93823 A1	21-01-2003
		US 6096918 A	01-08-2000
		ZA 9900567 A	26-07-1999
EP 0887338 A	30-12-1998	EP 0887335 A1	30-12-1998
		EP 0887338 A1	30-12-1998
		EP 1223160 A2	17-07-2002
		AU 725315 B2	12-10-2000
		AU 7309898 A	07-01-1999
		AU 722609 B2	10-08-2000
		AU 7309998 A	07-01-1999
		BR 9802199 A	20-07-1999
		BR 9802203 A	29-06-1999
		CA 2241391 A1	23-12-1998
		CA 2241397 A1	23-12-1998
		DE 69808557 D1	14-11-2002
		DE 69808557 T2	10-07-2003
		DE 69820928 D1	12-02-2004
		EP 0887336 A1	30-12-1998
		ES 2183313 T3	16-03-2003
		JP 11147852 A	02-06-1999
		JP 3124255 B2	15-01-2001
		JP 11116537 A	27-04-1999
		SG 93818 A1	21-01-2003
		US 6207857 B1	27-03-2001
		US 6232487 B1	15-05-2001
		ZA 9805262 A	11-01-1999
		ZA 9805265 A	11-01-1999

